## Serbian Biochemical Society

President: Marija Gavrović-Jankulović Vice-president: Suzana Jovanović-Šanta General Secretary: Isidora Protić-Rosić

Treasurer: Milica Popović

Scientific	Board

Marija GavrovićJankulović
Mihajlo B. Spasić
Vesna Niketić
Ivanka Karadžić
Svetlana Dinić
Nevena Đukić
Jelena Bašić
Ivan Spasojević

Ivana Beara Mojca Stojiljković Andjelka Ćelić Željko Popović

Žanka Bojić Trbojević

Milan Nikolić

Ana Ninić Adela Pitea

Zupkó István Vlatka Zoldos Aleksandra Inić-

Kanada

Tomasz Jurkowski Yaraslau Dzichenka Brankica Janković Sanja Krstić

Organization Committee

Suzana Jovanović-

Šanta

Jelena Purać

Milica Popović Emilija Svirčev Miloš Opačić

Milena Dimitrijević Tatjana Majkić Sofija Bekić Diandra Pintać Isidora Protić-Rosić Marina Crnković

Maja Marinović Iva Uzelac Jovana Drljača

Miloš Avramov Srđana Đorđievski Milana Bosanac

Vanja Tatić

#### **Proceedings**

Editor: Ivan Spasojević

Technical support: Jelena Korać Jačić

Cover design: Zoran Beloševac

Publisher: Faculty of Chemistry, Serbian Biochemical Society

Printed by: Colorgrafx, Belgrade

# Serbian Biochemical Society Eleventh Conference

Scientific meeting of an international character

September 22<sup>nd</sup> and 23<sup>rd</sup>, 2022, Novi Sad, Serbia

"Amazing Biochemistry"

### Antitumor properties of Eryngium amethystinum extract

Sanja Jelača<sup>1\*</sup>, Sanja Mijatović<sup>1</sup>, Zora Dajić-Stevanović<sup>2</sup>, Nebojša Arsenijević<sup>3</sup>, Danijela Maksimović-Ivanić<sup>1</sup>

Eryngium amethystinum is herb from Apiaceae family. It has been used as a food or in traditional medicine for the treatment of various diseases<sup>1</sup>. In the present work, we have evaluated its cytotoxic effect on a panel of human cancer cells (human breast carcinoma MCF7, human malignant melanoma A375, human colon carcinoma HCT116 and human lung cancer A549). Treatment with Eryngium amethystinum ethanol extract decreased viability of all cancer cell lines in a dose-dependent manner after 72 h. For further investigation of potential mechanism of action A549 cell line was selected. The observed viability decrease was followed by loss of dividing potential after the treatment. Aditionally, 90% of A549 cells were subjected to programmed cell death – apoptosis which was not followed with caspase activation. In parallel with this, typical apoptotic morphology of treated cells was observed by fluorescent microscopy. Apart from this decreased cell viability was due to triggered autophagic cell death. Taken together, the shown effect of Eryngium amethystinum makes this plant worthwhile for further evaluation in the field of oncology.

#### Acknowledgements

This study was supported by: Grant: Ministry of Education, Science and Technological development of the Republic of Serbia No. 451-03-9/2021-14/200007 and joint funding of Development and research projects of the Republic of Serbia and the People's Republic of China.

#### References

 Cianfaglione K, et al. Cytotoxic essential oils from Eryngium campestre and Eryngium amethystinum (Apiaceae) growing in central Italy. Chem Biodivers 2017;14: 10.1002/cbdv.201700096.

<sup>&</sup>lt;sup>1</sup>Institute for Biological Research "Siniša Stanković" – National Institute of Republic of Serbia, University of Belgrade, Belgrade, Serbia

<sup>&</sup>lt;sup>2</sup>Faculty of Agriculture, University of Belgrade

<sup>&</sup>lt;sup>3</sup>Center for Molecular Medicine and Stem Cell Research, Faculty of Medical Sciences, University of Kragujevac, Kragujevac, Serbia

<sup>\*</sup>e-mail: sanja.jelaca@ibiss.bg.ac.rs