



Snežana VUJIN¹
Valerija MATIĆ¹
Željko PRIJOVIĆ¹
Miroslav M. VRVIĆ²
Mihajlo B. SPASIĆ³

Vitamin E and serum lipid level in patients with rectal carcinoma

ABSTRACT

Background: Epidemiological studies indicated connection of the rectal carcinoma with the dietary lipid content, but the identification of the links associating the causes with the consequences are still far from being resolved. On the other hand, it has been experimentally confirmed that vitamin E expresses antitumour effects in some types of malignancies, but these effects strongly depend not only on the vitamin E form employed, but also on the presence of other antioxidants, e.g. selenium (Se).

Materials and methods: In this work, we monitored the values of the following parameters in blood of patients diagnosed as having rectal carcinoma: cholesterol, HDL, LDL, triglycerides and vitamin E. The measurements were performed immediately before and after the surgery (rectal resection), as well as 1, 3 and 6 months postoperatively. One group of patients was postoperatively supplemented with dietary antioxidants and the values of the above parameters were monitored by the same schedule.

Results: Ratio of vitamin E content and the sum of triglycerides and cholesterol (E/trg+chol) was lower in patients than in the population of adult healthy volunteers, suggesting it to be a more reliable indicator of the risk factor for rectal carcinoma development than the level of vitamin E alone. A decline of both vitamin E level and E/trg+chol ratio was noticed in patients as early as one month postoperatively, while in the group receiving antioxidants, a significant decrease of these parameters was recorded only 6 months postoperatively.

Conclusion: The fact that the supplementation postpones reduction of vitamin E level postoperatively suggests that antioxidants could express some beneficial effects on the outcome of a standard therapy.

Key words: Vitamin E; Triglycerides; Cholesterol; Rectal Cancer; Antioxidants

Archive of Oncology 2000,8(4):165-7©2000, Institute of Oncology Sremska Kamenica, Yugoslavia

¹INSTITUTE OF CHEMISTRY, TECHNOLOGY AND METALLURGY, BELGRADE, YUGOSLAVIA

²FACULTY OF CHEMISTRY, UNIVERSITY OF BELGRADE, BELGRADE, YUGOSLAVIA

³DEPARTMENT OF PHYSIOLOGY, INSTITUTE FOR BIOLOGICAL RESEARCH "SINIŠA STANKOVIĆ", BELGRADE, YUGOSLAVIA

INTRODUCTION

The data on the role of vitamin E in the etiology of the rectal carcinoma, as well as on the influence of a supplementation therapy on this disease, based on clinical experience, are rather scarce (1). Epidemiological studies indicated connection of the rectal carcinoma with the dietary lipid content, but the identification of the links associating the causes with the consequences are still far from being resolved (2). According to the data accumulated during the past several years, the diet consisting of moderate amounts of lipids and proteins and high

amounts of fruit and vegetables, rich in cellulose fibers, represents the primary measure in the prevention of the rectal carcinoma development (3). As a risk of rectal carcinoma is increased by mutagenic action of free radicals, dietary antioxidants such as vitamin E, through reduction of this harmful species should be protective in the case of rectal carcinoma. In the present study, relevant parameters of lipid profile cholesterol, triglycerides, HDL, LDL, as well as content of vitamin E as a liposoluble antioxidant were monitored. The objective of the protocol of this study was also to examine whether the increase of plasma of vitamin E due to supplementation improve lipid profile in patients with rectal carcinoma.

were obtained from Surgical Clinic Belgrade, and taken from patients of both sexes, (age range from 42 to 65 years). Sampling was done before and immediately after the resection of colon, as well as 1, 3 and 6 months after the surgical intervention. A group of 29 patients were supplemented with antioxidants while the remaining 13 patients served as the control. Supplementation started immediately after the surgical intervention, and blood serum samples were analyzed 1, 3 and 6 months later. Supplemented patients were given organically bound selenium in a dry, inactive biomass of beer yeast with vitamins A, C and E (Oligogal Se®, Galenika Pharmaceutical Works, Zemun Yugoslavia, (4)) 3x1 dose daily for the first 4 months of supplementation, and 1x1 daily beginning with the 5th month of treatment with recommendation to avoid fatty and salty food.

Address correspondence to:
Snežana Vujin, Institute of Chemistry, Technology and Metallurgy, Njegoševa 12, 11000 Belgrade, Yugoslavia

The manuscript was received: 15. 06. 2000.

Provisionally accepted: 04. 08. 2000.

Accepted for publication: 29. 09. 2000.

MATERIALS AND METHODS

Sample

Blood serums from 42 patients with rectal carcinoma were used in this work. Samples

Formulation of Oligogal Se®

Based on the results of determination of organically bound selenium in a dry, inactive



biomass of beer yeast enriched additionally with selenium ("selenium yeast"), a formulation Oligogal Se® was determined. A single dose consisted of organically bound selenium 100 µg in "selenium yeast", vitamin A 1500 IU, vitamin C 90 mg, and vitamin E 45 IU (4).

Determination of vitamin E content in blood serum

Vitamin E (R, R, R- α -tocopherol) in blood serum was determined by gas chromatography (5).

The exact volume of blood serum (ranged from 1 ml to 5 ml) was extracted with an equal volume 1:1 of hexane p.a. Merck 2-3 times in glass cuvettes. After extraction the samples were centrifuged at 3000-4000 rpm, 15-20 min. n-hexane extracts were separated and n-hexane evaporated up to the volume of 0.1 ml by nitrogen streaming. BHT (2,6-bi-terc-butyl-4-methyl phenol) was added to the samples and concentration of vitamin E was determined by Varian gas chromatograph, using super DB 5 column, 5 m long, with 0.33 mm radius.

The conditions of analyses: temperature of the detector was 300°C (flame ionization detector); temperature of the injector 200°C; temperature program and speed 50-285°C; 15°C/min; carrier gas - nitrogen; and -gases for combustion - oxygen, hydrogen.

Vitamin E concentration was calculated using the standard curve as reference. Normal values were from 15-24 mmol/L (4-6).

Determination of triglycerides, cholesterol, HDL and LDL

Triglycerides, cholesterol, HDL and LDL were determined by using standard biochemical procedures incorporated in Radox tests. The values up to 1.95 mmol/L for triglycerides, 3.1-6.5 mmol/L for cholesterol, 0.78-1.95 mmol/L for HDL and 1.55-4.53 mmol/L for LDL were taken as normal.

Statistical analyses

For statistical analyses of data standard Student's t-test was performed (7).

RESULTS

Our results show (Table 1) that in patients with rectal carcinoma the preoperative value of vitamin E ranges within normal scope of values. Ratio ($\mu\text{M}/\text{mM}$) of vitamin E content and the sum of triglycerides and cholesterol (E/trg+chol) was lower (R=2) in patients than in the population of adult healthy volunteers (R=4) (Figure 1), suggesting it to be a more reliable indi-

Table 1. Serum vitamin E, TG, CHL, HDL and LDL concentrations in patients before and after the colon resection - the effect of supplementation

	Non-supplemented			Oligogal Se®				
	Before	After	1	3	6	1	3	6
Vit E	15.96±3.14	13.60±3.10	1.52±1.26*	5.52±1.54*	2.65±1.50*	15.09±6.07	10.67±3.28	4.01±1.44*
TG	1.56±0.10	0.89±0.08	1.52±0.18	1.99±0.45	1.96±0.46	1.68±0.11	1.43±0.08	1.77±0.15
CHL	5.82±0.2	3.68±0.14	4.91±0.56	5.52±0.54	5.89±0.40	5.69±0.20	5.77±0.21	6.07±0.25
HDL	1.44±0.07	1.03±0.05	1.48±0.19	1.77±0.24	1.76±0.20	1.34±0.06	1.49±0.08	1.81±0.12
LDL	3.77±0.18	2.23±0.15	2.74±0.51	2.84±0.66	3.28±0.43	3.57±0.18	3.62±0.20	3.51±0.19

* p<0.005

Units for vitamin E are $\mu\text{mol}/\text{L}$, and for TG, CHL, HDL and LDL mmol/L. The results were expressed as means \pm S.E

cator of the risk factor for rectal carcinoma development than the level of vitamin E alone.

as *c-fos*, *c-jun* and *c-myc* as reported by several authors (11-14). On the other hand, reactive oxy-




Figure 1. The vitamin E/(TG+CHL) ratio in patients before and after the colon resection - the effect of supplementation. Ratio ($\mu\text{M}/\text{mM}$) for health volunteers was 4. The results were expressed as means \pm S.E. Statistical significance: * - p<0.005 in respect to group before or after the colon resection.

A decline of both vitamin E level and E/trg+chol ratio was noticed in non-supplemented patients as early as one month postoperatively, while in the group receiving antioxidants, a significant decrease of these parameters was recorded only 6 months postoperatively. No significant differences in the content of both HDL and LDL were observed neither when the patients were compared with healthy controls, nor when supplemented patients were compared with non-supplemented ones.

DISCUSSION

Constant oxidative stress in the induction of malignancies is significant from several aspects. First of all, a constant oxidative stress can permanently activate transcription factors and induce the expression of proto-oncogenes such

gen species (ROS) can damage specifically different protease inhibitors such as α_1 -proteinase inhibitor, α_2 -macroglobulin and α_2 -plasma inhibitor by oxidizing methionine residues in their active centers. Inactivation of these protease inhibitors affects the activities of the proteases, e.g. elastase and plasmin, thus accelerating tumour invasion and development of metastases (15). Recent studies revealed disturbed blood plasma oxidative status in patients with gastric carcinoma. Altered lipid peroxidation was accompanied by a significant decrease of vitamin E, ascorbate, β -carotene, urates and glutathione levels and an increased ceruloplasmin copper content (14).

Numerous studies were focused on the monitoring of antioxidants in body liquids because their antimutagenic and antitumour effects have been clearly demonstrated (16). It

has been shown that even vitamin E supplementation does not lead to restitution of its blood level in patients with rectal carcinoma, very probably because of the disturbance associated with the mechanism maintaining adequate vitamin E content in the cell membranes. Our results show that decline of both vitamin E level and E/trg+chol ratio occurred in patients as early as one month postoperatively, while in the group receiving antioxidants, a significant decrease of these parameters was recorded only 6 months postoperatively. In 1986, Traber et al. (17) observed a decreased efficiency of vitamin E absorption in the case of its increased intake, which can be of great importance when supplementation of this vitamin is recommended. In this study vitamin E was supplemented in daily dose of 130 U together with other antioxidants. It has been suggested that polyunsaturated long chain fatty acid-rich triglycerides and retinoic acid can lead to a significant decline in the extent of vitamin E absorption (17). Polyunsaturated long chain fatty acids were shown to accelerate vitamin E oxidation under *in vivo* conditions and the observed inhibition of its absorption can be of a special clinical interest (18).

As different antioxidants appear to act synergistically, supplementation with vitamin E might be more effective if combined with other micronutrients (19), as it was in our study. It is evident that this way of supplementation delays the postoperative decrease in vitamin E content and ratio VitE/ trg+chol thus reducing this risk factor (19).

REFERENCES

1. Stone WL, Papas AM. Tocopherol and etiology of colon cancer. *J Natl Cancer Inst* 1997;89:1006-14.
2. Wils J, Wagner DJ. Adjuvant treatment of colon cancer. Where to go from here. *Ann Oncol* 1990;1:329.
3. Vogelstein B. Genetic alteration during colorectal tumor development. *N Engl J Med* 1988;319:595.
4. Vrvic M, Vučetić J, Matic V, Veljković V, Lazić M. Biotechnological process for the production of the stable, dietetic product based on modified yeast enriched in selenium. Yugoslav Patent, 17.10.1988;No 46117.
5. Desai DI, Machlin LJ. Vitamin E. In: Klein A, Venugopal B, eds. *Methods of Vitamin Assay*. New York: John Wiley & Sons, 1985:255-83.
6. Machlin LJ, Gabriel E. Kinetics of tissue α -tocopherol uptake and depletion following administration of high levels of vitamin E. In: Lubin B, Machlin LJ, eds. *Vitamin E, biochemical, Hematological, and Clinical Aspects*. *Ann NY Acad Sci* 1982;393:48-59.
7. Sokal RR, Rohlf FJ, eds. *Biometry*. San Francisco: Freeman, 1981.
8. Peterlini NJ, Mittelman A. Colon and rectal carcinoma. In: *Handbook of chemotherapy and clinical oncology*. Paris: F. I. I. S. 1988;274-80.
9. Saito M, Kobatake Y, Tsuchida M, Kuroda K, Innami S, Itoh R. Nutritional status of vitamin E in a healthy elderly population in Japan. *J Clin Biochem Nutr* 1988;4:149-56.
10. Vanderwoude MFJ, Vanderwoude MG. Vitamin E status in a normal population: The influence of age. *J Am College Nutr* 1987;6:307-11.
11. Prasad KN. Mechanisms of action of vitamin E on mammalian tumor cells in culture. *Nutr Growth Canc* 1988;363-75.
12. Dreher D, Junod AF. Role of oxygen free radicals in cancer development. *Eur J Cancer* 1996;32A:1:30-8.
13. Kline K, Yu W, Zhao B, Israel K, Charpentier A, Menchaca MS et al. Vitamin E succinate: mechanisms of action as tumor cell growth inhibitor. In: Prasad KN, Santamaria L, Williams RM, eds. *Nutritients in cancer prevention and treatment*. Totowa, NY: Humana Press, 1995:39-56.
14. Miyamoto H, Araya Y, Ito M, Isobe H, Dosaka H, Shimizu T et al. Serum selenium and vitamin E concentration in families of lung cancer patients. *Cancer* 1987;60:1159-62.
15. Arvashagan S, Kavitha K, Nagini S. Plasma antioxidants in gastric carcinoma patients. *Med Sci Res* 1996;24:481-2.
16. Collins RH Jr. Colon cancer, dysplasia and surveillance in patients with ulcerative colitis. *N Engl J Med* 1987;316:1654-60.
17. Traber MG, Kayden HJ, Green JB. Absorption of water-miscible forms of vitamin E in a patients with cholestasis and in thoracic duct-cannulated rats. *Am J Clin Nutr* 1986;44:914-23.
18. Horwit MK. Interrelation between vitamin E and polyunsaturated fatty acids in adult men. *Vitamins and Hormones* 1962;20:541-58.
19. Pryor WA. Vitamin E and heart disease: basic science to clinical interventions trials. *Free Rad Biol Med* 2000;28:141-64.