

Македонски Фармацевтски Macedonian билтен Pharmaceutical Bulletin

волумен **68** (додаток 2) 2022 / volume **68** (Suppl 2) 2022



Македонско фармацевтско друштво, ул. Маршал Тито 13б/8, Скопје, Македонија

Macedonian Pharmaceutical Association, Marshal Tito 13b/8, Skopje Macedonia

Your hosts
Macedonian Pharmaceutical Association,
Faculty of Pharmacy, Ss ‘Cyril and Methodius’ University in Skopje, and
Association for Medicinal and Aromatic plants of Southeast European
Countries (AMAPSEEC)

Organizing committee

Gjoshe STEFKOV, Ss. Cyril and Methodius University in Skopje, Faculty of Pharmacy / Chair
Ivana CVETKOVIKJ KARANFILOVA, Ss. Cyril and Methodius University in Skopje, Faculty of Pharmacy / Secretary
Gabriela PETROVSKA-DIMITRIEVSKA, Ss. Cyril and Methodius University in Skopje, Faculty of Pharmacy / Registration officer
Olga GIGOPULU, Ss. Cyril and Methodius University in Skopje, Faculty of Pharmacy / Sponsorship officer
Ljubica MIHAILOVA, Ss. Cyril and Methodius University in Skopje, Faculty of Pharmacy / IT officer

Scientific committee

Biljana BAUER / Chair - North Macedonia	Sonja IVANOVSKA - North Macedonia
Zora DAJIĆ STEVANOVIĆ - Serbia	Nada KOVAČEVIĆ - Serbia
Svetlana KULEVANOVA - North Macedonia	Ufuk KOCA CALISKAN - Turkey
Vassya BANKOVA - Bulgaria	Marija JUG-DUJAKOVIĆ - Croatia
Nazim SEKEROGLU - Turkey	Krystyna SKALICKA-WOZNIAK - Poland
Dea BARIČEVIČ - Slovenia	Dejan PLJEVLJAKUSIĆ - Serbia
Kalina ALIPIEVA - Bulgaria	Maria HALABALAKI – Greece
Gjoshe STEFKOV - North Macedonia	Alban IBRALIU – Albania
Dragoja RADANOVIĆ - Serbia	Ye YANG – China
Katarina SAVIKIN - Serbia	Ivan SALAMON - Slovakia
Gordana POPSIMONOVA - North Macedonia	Antoaneta TRENDABILOVA - Bulgaria
Ilkay ERDOGAN ORHAN - Turkey	Biljana MIOVA - North Macedonia
Antonios MAKRIS - Greece	Lijiang XUAN - China
Marija KARAPANDZOVA -North Macedonia	Zlatko ŠATOVIĆ - Croatia
Xin XIE - China	Thomas JAKSCHITZ - Austria
Milen I. GEORGIEV – Bulgaria	Slavco HRISTOVSKI - North Macedonia
Sonja GADZOVSKA SIMIC - North Macedonia	Reneta GEVRENOVA - Bulgaria
Ivana CVETKOVIKJ KARANFILOVA - North Macedonia	Éva ZAMBORINE NEMETH - Hungary
	Avni HAJDARI – Kosovo

This issue of *Macedonian Pharmaceutical Bulletin* contains short communications accepted by the Scientific Committee for the presentation at the 11th Conference of Medicinal and Aromatic Plants of Southeast European Countries 2022 (CMAPSEEC 2022).

The authors are fully responsible for the contents of their short communications.

All reviewers that were involved in the short communications revision process are sincerely acknowledged.

Anti-melanoma effects of ingenanes isolated from *Euphorbia* species

Gordana Krstić^{*1}, Milka Jadranin², Sofija Jovanović Stojanov³, Milica Pešić³,
Vele Tešević¹, Slobodan Milosavljević^{1,4}

¹University of Belgrade – Faculty of Chemistry, Studentski trg 12–16, 11000 Belgrade, Serbia

²University of Belgrade – Institute of Chemistry, Technology, and Metallurgy, Department of Chemistry,
Njegoševa 12, 11000 Belgrade, Serbia

³Department of Neurobiology, Institute for Biological Research “Siniša Stanković” – National Institute of the Republic
of Serbia, University of Belgrade, Bulevar despota Stefana 142, 11060 Belgrade, Serbia

⁴Serbian Academy of Sciences and Arts, Kneza Mihaila 35, 11000 Belgrade, Serbia

Introduction

The Euphorbiaceae family is one of the most widespread plant families containing about 300 genera and more than 8000 species (Webster, 1994). The plants belonging to this genus are well known for the chemical diversity of their terpene constituents. Literature data indicate that metabolites isolated from this genus have analgesic, antipyretic, antimicrobial, antiviral, antiproliferative, and selective antitumor activity (Jadranin et al., 2013, Krstić et al., 2018, Wang et al., 2003). The most important metabolite isolated from the genus is ingenol-mebutate. It was isolated from the latex of *Euphorbia paralias* and *E. peplus*, and it has been registered under phase III clinical trials under the commercial name Picato® for the treatment of actinic keratosis and pre-cancerous and cancerous skin changes (Aditya and Gupta, 2013, Seca and Pinto, 2018).

In this research, from two species, *E. palustris* and *E. lucida*, four ingenane derivatives were isolated. Their anticancer effects were evaluated in the human melanoma – 518A2 cell line and compared with the effects of ingenol-mebutate. Selectivity towards human melanoma cells was determined using normal human keratinocytes – HaCaT.

Materials and methods

Plant materials

The *E. palustris* latex was collected in Besni Fok (Serbia), in May 2011, while the *E. lucida* stems were collected in Makiš (Serbia), in May 2012. The plants were identified by Professor Petar Marin, University of Belgrade – Faculty of Biology, Institute of Botany. Voucher specimens (No. 16,877 for *E. palustris*, and No. 16,879 for *E. lucida*) have been deposited at the Herbarium of Botanical Garden “Jevremovac”, University of Belgrade, Belgrade (Serbia).

Isolation and purification

The collected latex of *E. palustris* was first lyophilized and then the obtained lyophilizate was separated by gradient *dry-flash* column chromatography (DF CC). Fractions obtained by elution with 60 and 80% EtOAc were further fractionated by open column chromatography. Selected fractions were finally purified on RP HPLC. From these fractions, ingenanes **1** (9.0 mg) and **2** (6.1 mg) were isolated. *E. lucida* stems were dried and ground and then extracted with EtOAc/hexane mixture. The resulting extract was further fractionated with DF CC, gradient from 100% hexane to 100% EtOAc. Fractions eluted with 15, 40, and 60% EtOAc were further separated first by column chromatography, and RP HPLC was used for final purification. From these fractions, ingenanes **1** (5.5 mg), **3** (2.2 mg), and **4** (7.5 mg) were isolated.

Cell lines and assays

518A2 and HaCaT cells were cultured in DMEM supplemented with 10% FBS, 4 g/L glucose, L-glutamine (2 mM), 5000 U/mL penicillin, and 5 mg/mL streptomycin solution. All cell lines were sub-cultured at 72 h intervals using 0.25% trypsin/EDTA and seeded into a fresh medium at the density of 8000 cells/cm².

Cell viability was assessed by MTT assay. To determine IC₅₀ values of ingenol-mebutate, **1**, **2**, **3**, and **4**, cells were treated with different increasing concentrations of compounds. The absorbance of obtained dye was measured at 570 nm with a reference wavelength of 690 nm. IC₅₀ was calculated by non-linear regression analysis.

The percentages of apoptotic, necrotic, and viable cells were assessed by Cell Death Detection Kit and flow cytometry using Annexin-V-FITC and PI staining according to the manufacturer's instructions.

Results and discussion

Ingenanes **1** (3 β -benzoyloxy-13 α -dodecanoyloxy-ingenol) (Wang et al., 2003) and **2** (3 β ,13 α ,17-tribenzoyloxy-ingenol) (Lu et al., 2008) were isolated from *E. palustris* latex, while a total of three ingenane derivatives were isolated from *E. lucida* stem: **1**, **3** (3 β ,5 β -dibenzoyloxy-20-deoxyingenol) (Gotta et al., 1984) and **4** (3 β ,5 β ,20-tribenzoyloxyingenol) (Opferkuch et al., 1981). All isolated compounds have been described in the literature, but compound **4** was reported as a natural product for the first time. Previously, compound **4** has been described in the literature as the esterification reaction product of the ingenol. The structures of the isolated compounds were determined using data obtained from 1D and 2D NMR spectra, and the molecular formulas were confirmed by HR-ESI MS. In the literature, the biological activities of the isolated compounds have not been described in detail.

Ingenanes **1** and **2** exerted a strong cytotoxic effect in the nanomolar range (IC₅₀ values were 5 nM and 45 nM, respectively) in melanoma cells, while normal keratinocytes were not sensitive to these compounds even in the micromolar range. Ingenol-mebutate, **3**, and **4** had a less prominent anti-melanoma effect with IC₅₀ values between 20 μ M and 30 μ M in 518A2 cells. Selectivity towards melanoma cells was considerable for ingenol-mebutate and **4** with IC₅₀ values in HaCaT cells > 100 μ M. Cell Death analysis confirmed the preferential cytotoxicity of ingenanes in melanoma cells.

Conclusion

Ingenanes isolated from *E. palustris* and *E. lucida*

showed significant and selective anti-melanoma effects and further molecular mechanism elucidation *in vitro* and *in vivo* is envisioned.

Acknowledgment

This work was financially supported by the Serbian Academy of Sciences and Arts (Strategic Project MilkIng, No. 01-2022), and by Ministry of Education, Science and Technological Development of the Republic of Serbia (Contract Nos: 451-03-68/2022-14/200168, 451-03-68/2022-14/200026 and 451-03-68/2022-14/200007).

References

- Aditya, S., Gupta, S., 2013. Ingenol mebutate: A novel topical drug for actinic keratosis. *Indian Dermatol. Online J.* 4(3), 246-249. <https://doi.org/10.4103/2229-5178.115538>.
- Gotta, H., Adolf, W., Opferkuch, H.J., Hecker, E., 1984. On the active principles of the Euphorbiaceae, IXa ingenane type diterpene esters from five *Euphorbia* species. *Z. Naturforsch., B: J. Chem. Sci.* 39(5), 683-694. <https://doi.org/10.1515/znb-1984-0525>.
- Jadranin, M., Pešić, M., Aljančić, I.S., Milosavljević, S.M., Todorović, N.M., Podolski-Renić, A., Banković, J., Tanić, N., Marković, I., Vajs, V.E., Tešević, V.V., 2013. Jatrophone diterpenoids from the latex of *Euphorbia dendroides* and their anti-P-glycoprotein activity in human multi-drug resistant cancer cell lines. *Phytochemistry.* 86, 208-217. <https://doi.org/10.1016/j.phytochem.2012.09.003>.
- Krstić, G., Jadranin, M., Todorović, N.M., Pešić, M., Stanković, T., Aljančić, I.S., Tešević, V.V., 2018. Jatrophone diterpenoids with multidrug-resistance modulating activity from the latex of *Euphorbia nicaeensis*. *Phytochemistry.* 148, 104-112. <https://doi.org/10.1016/j.phytochem.2018.01.016>.
- Lu, Z.Q., Yang, M., Zhang, J.Q., Chen, G.T., Huang, H.L., Guan, S.H., Ma, C., Liu, X., Guo, D.A., 2008. Ingenane diterpenoids from *Euphorbia esula*. *Phytochemistry.* 69(3), 812-819. <https://doi.org/10.1016/j.phytochem.2007.09.013>.
- Opferkuch, H.J., Adolf, W., Sorg, B., Kusumoto, S., Hecker, E., 1981. On the chemistry of ingenol, I ingenol and some of its derivatives. *Z. Naturforsch., B: J. Chem. Sci.* 36(7), 878-887. <https://doi.org/10.1515/znb-1981-0717>.
- Seca, A.M.L., Pinto, D.C.G.A., 2018. Plant Secondary Metabolites as Anticancer Agents: Successes in Clinical Trials and Therapeutic Application. *Int. J. Mol. Sci.* 19(1), 263. <https://doi.org/10.3390/ijms19010263>.
- Wang, L.Y., Wang, N.L., Yao, X.S., Miyata, S., Kitanaka, S., 2003. Diterpenes from the roots of *Euphorbia kansui* and their in vitro effects on the cell division of *Xenopus*. *Chem. Pharm. Bull.* 51(8), 935-941. <https://doi.org/10.1248/cpb.51.935>.
- Webster, G.L., 1994. Classification of the Euphorbiaceae. *Ann. Missouri Bot. Gard.* 81(1), 3-32. <https://doi.org/10.2307/2399908>.